

# Optimization

## Problem set 1

Due Monday, April 12th

1. Consider the closed convex set  $B_1 = \{x \in \mathbb{R}^n \mid \|x\|_1 = \sum_i |x_i| \leq 1\}$ . This is the unit ball of the  $\ell_1$  norm.
  - (a) Show that  $B_1$  is a polyhedron by explicitly expressing it as an intersection of halfspaces. How many halfspaces (“facets”) are required in order to express  $B_1$ ?
  - (b) Explicitly express  $B_1$  as a convex hull of a finite number of points. How many points (“vertices”) are required in this characterization?
  - (c) Contrast this with the  $\ell_\infty$  unit ball,  $B_\infty = \{x \in \mathbb{R}^n \mid \|x\|_\infty \leq 1\}$ . How many halfspaces are required in order to express  $B_\infty$  as an intersection of halfspaces? How many points are required in order to express  $B_\infty$  as a convex hull?
  - (d) For each point  $\hat{x}$  on the boundary of  $B_1$ , identify the set of all supporting hyperplanes of  $B_1$  at  $\hat{x}$  explicitly. For each such  $\hat{x}$ , what is the dimensionality of this set?
2. Consider a polyhedron  $C = \text{conv}\{v_1, \dots, v_k\} \subset \mathbb{R}^n$  and a convex function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ .
  - (a) Prove that a maximum of  $f$  over  $C$  is achieved at one of the vertices  $v_i$ . (Hint: assume the statement is false and use Jensen’s inequality). Is it possible that the maximum is also achieved at an interior point?

(A generalization of the above is that a maximum of a function over a closed and bounded convex set is achieved at an extreme point, i.e. a point which is not a convex combination of other points in the set).
  - (b) Use the above to conclude that the *minimum* of a linear objective over the polyhedron  $C$  is always achieved at one of the vertices  $v_i$ .

3. In this problem we will define strong convexity more generally than it is defined by Boyd and Vandenberghe (Section 9.1.2). In particular, we will consider a definition that is valid also for non-differentiable functions.

**Definition:** A function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is  $m$ -strongly convex if for every  $x, y \in \mathbb{R}^n$  and every  $\theta \in [0, 1]$ :

$$f((1 - \theta)x + \theta y) \leq (1 - \theta)f(x) + \theta f(y) - \frac{m}{2}\theta(1 - \theta) \|x - y\|_2^2$$

- (a) Prove that a continuously differentiable function  $f$  is  $m$ -strongly convex if and only if for every  $x, y \in \mathbb{R}^n$ ,

$$f(y) \geq f(x) + \nabla f(x)^T(y - x) + \frac{m}{2} \|y - x\|_2^2.$$

This generalizes the first order characterization of convexity (Section 3.1.3).

- (b) Prove that a twice continuously differentiable function  $f$  is  $m$ -strongly convex if and only if its domain is convex and for every  $x \in \mathbb{R}^n$ , all eigenvalues of the Hessian at  $x$  are greater or equal to  $m$ , i.e.:

$$\nabla^2 f(x) \succeq mI.$$

This generalizes the second order characterization of convexity (Section 3.1.4) and is the definition used in Section 9.1.2.

- (c) Provide an example of a function that is strongly convex but not everywhere differentiable.
- (d) Let  $f$  be a  $m$ -strongly convex function, and  $x^*$  an optimum for  $\min_{x \in \mathbb{R}^n} f(x)$ . Prove that for any point  $x \in \mathbb{R}^n$ :

$$f(x) \geq f(x^*) + \frac{m}{2} \|x - x^*\|_2^2.$$

Conclude that the optimum is unique and that any  $\epsilon$ -suboptimal point must be close to the optimum. Provide an explicit upper bound on  $\|x - x^*\|_2$  for an  $\epsilon$ -suboptimal  $x$ . (Note that if  $f$  is convex but not strongly convex,  $\epsilon$ -suboptimal points can be arbitrarily far away from the closest optimum).

Recommended review exercises from Boyd and Vandenberghe (please do not turn these in—they will *not* be graded): 2.12, 2.15, 3.6, 3.16, 3.18, 3.24, 3.26.